




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

SARS-CoV-2 transmission risk upon return to work in RNA-positive healthcare workers

L.M. Kolodziej, S. Hordijk, J. Koopsen, J.J. Maas, H.T. Thung, I.J.B. Spijkerman, M. Jonges, M.K. Bomers, J.J. Sikkens, M.D. de Jong, R. Zonneveld, J. Schinkel



PII: S0195-6701(22)00079-2

DOI: <https://doi.org/10.1016/j.jhin.2022.02.024>

Reference: YJHIN 6625

To appear in: *Journal of Hospital Infection*

Received Date: 28 November 2021

Revised Date: 21 February 2022

Accepted Date: 24 February 2022

Please cite this article as: Kolodziej LM, Hordijk S, Koopsen J, Maas JJ, Thung HT, Spijkerman IJB, Jonges M, Bomers MK, Sikkens JJ, Jong d, Zonneveld R, Schinkel J, SARS-CoV-2 transmission risk upon return to work in RNA-positive healthcare workersSummry, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2022.02.024>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 The Author(s). Published by Elsevier Ltd on behalf of The Healthcare Infection Society.

SARS-CoV-2 transmission risk upon return to work in RNA-positive healthcare workers

L.M. Kolodziej^{a*,†}, S. Hordijk^{a*}, J. Koopsen^a, J.J. Maas^b, H.T. Thung^b, I.J.B. Spijkerman^a, M. Jonges^a, M.K. Bomers^c, J.J. Sikkens^c, M.D. de Jong^a, R. Zonneveld^{a‡}, J. Schinkel^{a‡}

^a Department of Medical Microbiology and Infection Prevention, Amsterdam University Medical Centres, University of Amsterdam, Amsterdam, The Netherlands

^b Department of Occupational Health, Amsterdam University Medical Centres, University of Amsterdam, Amsterdam, The Netherlands

^c Department of Internal Medicine, Division of Infectious Diseases, Amsterdam University Medical Centres, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

^{*,‡} Authors contributed equally

[†] **Corresponding Author:** Lisa Marie Kolodziej

Department of Medical Microbiology and Infection Prevention, Amsterdam University

Medical Centres, University of Amsterdam

Meibergdreef 9, 1105 AZ Amsterdam, the Netherlands

Email: l.m.kolodziej@amsterdamumc.nl

Telephone: + 316 19809909

Running title: SARS-CoV-2 transmission among healthcare workers

Key words: SARS-CoV-2; COVID-19; Healthcare workers; Infectious disease transmission; Infection prevention and control

Declarations of interest: none

Word count (excluding summary and references): 2497

SUMMARY

Background

Healthcare workers (HCWs) are at risk for coronavirus disease 2019 (COVID-19) and for spreading Severe Acute Respiratory Syndrome Virus 2 (SARS-CoV-2) amongst colleagues and patients.

Aim

We aimed to study presence of SARS-CoV-2 RNA and possible onward transmission by HCWs upon return to work after COVID-19, and association with disease severity and development of antibodies over time.

Methods

Unvaccinated HCWs with positive SARS-CoV-2 RT-PCR were prospectively recruited. Data on symptoms was collected via telephone questionnaires on day 2, 7, 14 and 21 after positive test. Upon return to work, repeat SARS-CoV-2 RT-PCR was performed and serum was collected. Repeat sera were collected at week 4, 8, 12 and 16 to determine antibody dynamics over time. Phylogenetic analysis was conducted to investigate possible transmission events originating from HCW with a positive repeat RT-PCR.

Findings

Sixty-one (84.7%) participants with mild-moderate COVID-19 had a repeat SARS-CoV-2 PCR performed upon return to work (median 13 days post symptom onset), of which 30 (49.1%) were positive with a median cycle threshold (Ct) value of 29.2 (IQR 3.0). All HCWs developed antibodies against SARS-CoV-2. No significant differences in symptomatology and presence of antibodies were found between repeat RT-PCR-positive and -negative HCWs. Eleven direct colleagues of six participants with a repeat RT-PCR Ct-value <30 tested positive after the HCW returned to work. Phylogenetic and epidemiologic analysis did not

indicate onward transmission through HCW who were SARS-CoV-2 RNA positive upon return to work.

Conclusions

HCWs regularly return to work with substantial SARS-CoV-2 RNA loads. However, we found no evidence for subsequent in-hospital transmission.

Key words: SARS-CoV-2; COVID-19; Healthcare worker; Infectious disease transmission

INTRODUCTION

Healthcare workers (HCWs) play a critical role in the response against the ongoing coronavirus disease 2019 (COVID-19) pandemic. Multiple studies show higher infection rates in HCWs compared to the general population, suggesting an occupational risk.[1-3] As for all confirmed cases, COVID-19 in HCWs requires measures to prevent transmission including quarantine. Hereby, (long) periods of absence can increase the strain on the healthcare system. During this study, hospital guidelines prescribed that HCWs with confirmed COVID-19 could return to work 24 hours post symptom resolution. National and international guidelines generally recommend a minimal duration of isolation of 7 to 10 days after onset of COVID-19 symptoms and 24 hours to 5 days after improvement or resolution of symptoms.[4-7] Some guidelines mention the option of re-testing before returning to work for specific occasions (e.g., for HCWs with severe immune deficiencies),[5, 6, 8] but standard re-testing before returning to work is not recommended by any of the other guidelines since the assumed risk of transmission is considered negligible after these time periods.[9, 10]

On the other hand, Severe Acute Respiratory Syndrome Virus 2 (SARS-CoV-2) RNA can be detected in upper respiratory tract samples for prolonged periods, even without symptoms.[11] These cases are considered not to be infectious, as studies in mild cases of COVID-19 have found that no viable virus could be detected in individuals with prolonged shedding of SARS-CoV-2 RNA.[12, 13] However, in these studies samples were collected from 14 up to 30 days after diagnosis, whereas most HCWs may resume work sooner. In addition, in these studies viral culture was performed to determine infectivity and corresponding transmission risk. Since the standard procedure for HCWs returning to work in Dutch hospitals after a SARS-CoV-2 infection does not include RT-PCR or viral culture, viral loads at that time are not determined and the risk of transmission by mild cases who may return to work sooner remains unclear.

Repeat RT-PCR testing could further examine the risk of transmission of HCWs upon return to work. Furthermore, the presence of SARS-CoV-2 specific antibodies has been negatively correlated with the presence of infectious virus.[14, 15] Therefore, antibody dynamics could be valuable in determining the risk of transmission upon return to work and subsequent re-infection in this population with an increased occupational risk.

The aim of this prospective observational study is to assess the presence of SARS-CoV-2 RNA and corresponding cycle threshold (Ct) values upon resolution of symptoms in SARS-CoV-2 infected HCWs and its relation to disease severity, antibody dynamics and the risk of transmission.

METHODS

Study design

Participants

The Amsterdam University Medical Centres (Amsterdam UMC), the Netherlands, offers SARS-CoV-2 RT-PCR testing of combined nasopharyngeal and oropharyngeal swab specimens for HCWs with COVID-19-like symptoms (coughing, pharyngitis, dyspnoea, rhinitis and anosmia or dysgeusia). HCWs that tested positive in routine testing between May and September 2020, during the national ‘second wave’ and before the national vaccination campaign started, were invited to participate in this prospective observational study.

Sampling process

At day 2 after the positive SARS-CoV-2 RT-PCR, a telephone questionnaire regarding signs and symptoms at the time of disease onset as well as at the present time was administered. Hereby the presence of 14 predefined symptoms (coughing, pharyngitis, dyspnoea, rhinitis, abdominal pain, diarrhoea, nausea, vomiting, anorexia, fever, myalgia, headache, fatigue and

anosmia or dysgeusia) was determined. Follow-up symptomatology questionnaires were conducted at day 7, 14 and 21, as long as participants reported to experience symptoms.

Repeat nasopharyngeal and oropharyngeal swabs and initial serum were collected when HCWs returned to work. Hospital guidelines for returning to work required that all respiratory symptoms had to be resolved > 24 hours. Anosmia, dysgeusia and fatigue were not required to be resolved upon return to work. Repeat sera were collected at week 4, 8, 12 and 16 after the initial positive RT-PCR. All sera were stored at -20°C until serological tests were performed.

The nasopharyngeal and oropharyngeal swabs were collected in E-swab or UTM viral transport medium (COPAN Diagnostics, Murrieta, CA, USA).

Laboratory assays

SARS-CoV-2 RNA was extracted using the MagNA Pure 96 system (Roche, Penzberg, Germany). RT-PCR targeting the SARS-CoV-2 E gene was performed according to a previously published protocol.[16] The presence of antibodies was determined by the ELISA-based Wantai SARS-CoV-2 double antigen sandwich total antibody assay (Wantai Biological Pharmacy, Beijing, China).

Contact tracing in HCW that returned to work

Standard contact tracing was performed for every SARS-CoV-2 positive HCW (or patient) by the Infection Control department. To investigate the transmission risk of HCWs with a positive repeat PCR, potential secondary infections were identified using data of the Occupational Health and Infection Control department. Potential secondary infections were defined as contacts within the same department that tested positive for SARS-CoV-2 within 7 days after study participants with a repeat RT-PCR Ct-value <30 returned to work.

Viral genomes of specimens of study participants and return-to-work contacts were amplified using the Ion AmpliSeq™ SARS-CoV-2 Research Panel and sequenced on an Ion GeneStudio S5 system (both from ThermoFisher Scientific, The Netherlands). Sequences were phylogenetically analysed to infer relatedness in a background of contemporaneous SARS-CoV-2 viral genomes from the Netherlands, derived from the GISAID database (Table S1). A maximum-likelihood phylogeny was constructed using the Augur pipeline.[17] We used procedures taken from [github.com/nextstrain/ncov] including the clock rate, reference genome, and site masking. Trees were visualised using ggtree[18] as implemented in R (R Core Team, Vienna, Austria).

Ethics and Consent

Informed consent was obtained from all participants. The study was reviewed and approved by the Amsterdam UMC institutional review board and conducted in accordance with the Declaration of Helsinki, and national and institutional standards.

Statistical Analysis

Unknown or missing answers in the symptomatology questionnaires were considered as absent. Fatigue and anosmia/dysgeusia were not included to determine disease duration. Sera with an absorbance/cut off ratio (s/c) above 1.1 were considered positive, samples with an s/c below 0.9 were considered negative. A s/c between 0.9 and 1.1 was considered indeterminate.

The data was analysed using RStudio (R Core Team, Vienna, Austria) and Graphpad Prism version 9.0.2 for Mac (GraphPad Software, San Diego, California USA). Normality checks were performed using the Shapiro-Wilk test. Descriptive analyses were made on baseline characteristics and the number of observations, presented as numbers and percentages. For

descriptive statistics, quantitative variables that did not follow a normal distribution were presented with median and interquartile range (IQR). Binomial logistic regression was used to calculate odds ratios and 95% CI for evaluating the association of the presence of symptoms with seroprevalence and presence of viral RNA. P values <0.05 were considered significant.

RESULTS

Participants

A total of 72 HCWs were included in this study. Demographics are shown in Table I. One HCW was admitted to the hospital (1.4%). Upon study inclusion, 20.8% of the HCWs reported to have worked while having COVID-like symptoms before they tested positive. Experiencing mild symptoms that were not directly recognized was the most common explanation.

Symptomatology

The median time between disease onset and time of initial RT-PCR was 1 day (range 1-7). The median duration of symptoms was 10 days (range 0-41). Symptoms decreased over time (Table II). Fever and dyspnoea were not frequently reported. At disease onset, rhinitis, headache and fatigue were most frequently observed. Gastro-intestinal symptoms were reported in a minority of the HCWs. At day 21, 43% still reported symptoms. Fatigue and anosmia or dysgeusia most frequently persisted at day 21. The majority (80.6%) of HCWs had a self-reported mild experience of COVID-19. No significant differences in symptomatology were found between repeat RT-PCR-positive and repeat RT-PCR-negative HCWs (data not shown).

Virology

The median Ct-value of the initial RT-PCR was 21.1 (IQR 8.0). Sixty-one (84.7%) participants had a repeat RT-PCR performed upon return to work, with a median of 13 days (range 6-42) post symptom onset. Thirty (49.1%) of them were positive with a median Ct-value of 29.2 (IQR 3.0). Eleven participants did not have a repeat RT-PCR performed.

Twenty-two out of the 30 repeat RT-PCR-positive participants (73.3%) had a repeat RT-PCR specimen with a Ct-value <30 (corresponding with 36% of all HCW for which repeat RT-PCR results were available). Of these 22 participants, we identified eleven SARS-CoV2 RNA-positive within-department-contacts as potential secondary transmissions. Specimens of these eleven within-department-contacts were sequenced (Figure 1).

Phylogenetic analysis revealed one pair of identical viral genomes of return-to-work and corresponding within-department-contact and one pair that differed two single-nucleotide polymorphisms. Contact tracing and epidemiological data of these two pairs showed no indications of onward transmission. Eight return-to-work and corresponding within-department-contact pairs had pairwise genetic distances not compatible with direct transmission (minimal pairwise genetic distance of five single-nucleotide polymorphisms).

Serology

All HCWs of which serum was collected developed antibodies during the follow-up period (data not shown). Upon symptom resolution, antibodies were detected in 42 out of 48 (87.5%) HCWs of which serum was collected at this time point. At 16 weeks, antibodies were detected in 97.5% of the HCWs. Two HCWs seroreverted (from positive to negative antibody status) during the follow-up period, within 8 weeks after disease onset. No significant difference in presence of antibodies was found between repeat RT-PCR-positive and repeat RT-PCR-negative HCWs.

DISCUSSION

HCWs are at increased risk for SARS-CoV-2 infection and onward transmission to colleagues and patients. Guidelines are inconsistent on the timing for SARS-CoV-2 positive HCWs to return to work. We studied symptoms, repeated RT-PCR, risk of transmission and antibody dynamics in HCWs when returning to work. We found a generally mild course of COVID-19 and despite high SARS-CoV-2 RNA viral loads, no evidence for transmission from returning HCWs upon resolution of symptoms was found.

Surprisingly, almost 50% of the repeat RT-PCR when returning to work were positive with Ct-values suggesting the possibility of replicating virus. Our study showed RT-PCR positivity up to 38 days after symptom onset, which is in line with the now well-established experience that RNA may be detected for longer periods after a SARS-CoV-2 infection.[9-11] The relatively high viral loads (Ct-values <30) found in 36% of the HCW upon return to work in our study raised the question whether our hospital guideline is stringent enough to prevent nosocomial transmission, especially since national and international guidelines generally recommend a longer duration of isolation after COVID-19 in HCWs.[4-7]

Ct-values were used as surrogate marker for infectivity in accordance with previous studies, as they correlate well with the ability to culture (viable) virus and a cut-off of 30 is associated with the inability to culture virus. [19,20] Viral sequencing was performed to investigate whether onward transmission occurred by HCW who returned to work. Phylogenetic analysis showed one pair of identical viral sequences of a return-to-work study participant and within-department-contact and one pair that differed two single-nucleotide polymorphisms. For the pair with identical sequences the probability of direct transmission was deemed negligible after assessment of the contact tracing data as the index HCW worked from home during one

month after his infection and there was no contact to other HCWs at that time.

Epidemiological assessment of the pair differing two single-nucleotide polymorphisms suggested that direct transmission was unlikely, as the return-to-work HCW remained home for 14 days after onset of complaints, had no complaints when returning to work, and the HCWs did not know each other. Thus, despite the high numbers of positive specimens with theoretically viable virus in this study, we found no evidence for onward transmission at work from returning HCW upon resolution of symptoms. However, the possibility of HCW-to-HCW transmission cannot be completely ruled out as in this study onward transmission may have occurred but remained undiagnosed in asymptomatic individuals.

A possible explanation for the identical viral genomes found in one return-to-work and corresponding within-department-contact pair may be exposure to comparable genomes circulating in The Netherlands at that time (as evidenced by identical genomes detected in contemporaneous SARS-CoV-2 viral genomes from the Netherlands, Table A.I). Although direct transmission could not definitely be ruled out for one pair in this study, a symptom-based strategy for determining when HCWs with a SARS-CoV-2 infection could return to work as in the current hospital guidelines are considered adequate and safe. Nevertheless, as this study was performed before the emergence of the alpha-variant, the emergence of new circulating variants associated with higher transmissibility[21, 22] may require guideline re-evaluation. Moreover, as study participation was on a voluntary basis, the included HCW population may have behaved more compliant with social distancing rules and personal protection guidelines. This could partially explain the absence of documented transmission by HCW after returning to work. Infection prevention measures such as physical distancing, personal protective equipment and vaccination should remain a priority for SARS-CoV-2 in-hospital infection control, as there is evidence that HCW-to-HCW transmission is an

important route of nosocomial infections[22-25] and transmissions generally occur before a HCW tests positive.

Despite low symptomatology, all HCWs in this cohort seroconverted. Comparable prospective studies showed similar but somewhat lower rates, possibly due to a shorter follow up period[26, 27] or because only IgG was measured.[28] Further research is needed to determine long-term protection and protection against new variants. Presence of antibodies seemed not associated with repeat RT-PCR positivity, indicating that even mild infections with a faster viral clearance result in antibody response. The majority of the participants (87.5%) had already developed antibodies when returning to work, which further reduces the assumed risk of transmission at this time point given the negative correlation with SARS-CoV-2 specific antibodies and the presence of infectious virus.[14, 15]

The main limitation of our study is that infectivity of the HCWs when returning to work could not be determined. In addition, the small sample size of our study, especially the limited number of HCWs returning to work with high viral loads, may have influenced our conclusions about the risk of transmission. However, extensive phylogenetic as well as background analyses in combination with contact tracing data showed no evidence for direct transmission.

A strength of this study is that it was prospectively conducted in confirmed SARS-CoV-2 positive HCWs. Most studies in HCWs are retrospective seroprevalence studies in which it is impossible to accurately evaluate symptomatology or determine the antibody responses in this specific population. Furthermore, all analyses were performed in the same laboratory, making it possible to compare Ct-values amongst participants.

Conclusions

269 To conclude, our study revealed relatively high viral loads in SARS-CoV-2 positive HCWs
270 when returning to work after symptom resolution. As no evidence for secondary HCW-to-
271 HCW transmission after returning to work was found, a symptom-based approach appears
272 adequate in preventing SARS-CoV-2 infections from returning HCW. Since HCW-to-HCW
273 transmission is a common source of nosocomial SARS-CoV-2 infections, infection prevention
274 measures and guideline adherence should remain priorities when shaping future hospital
275 policy and practice.

ACKNOWLEDGEMENTS

We give special thanks to the laboratory team including dr. Matthijs Welkers, dr. Robin van Houdt, dr. Rosa van Mansfeld and Sjoerd Rebers for conducting the (sequencing) analyses and to the team of the department of Occupational Health for providing information to the (possible) participants and obtaining the specimens.

DECLARATIONS OF INTEREST

None.

FUNDING STATEMENT

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

288 REFERENCES

- 289 1. Nguyen LH, Drew DA, Graham MS, et al. Risk of COVID-19 among front-line
290 health-care workers and the general community: a prospective cohort study. *Lancet Public*
291 *Health*. 2020;**5**(9):e475-e83.
- 292 2. Rudberg AS, Havervall S, Manberg A, et al. SARS-CoV-2 exposure, symptoms and
293 seroprevalence in healthcare workers in Sweden. *Nat Commun*. 2020;**11**(1):5064.
- 294 3. Mutambudzi M, Niedwiedz C, Macdonald EB, et al. Occupation and risk of severe
295 COVID-19: prospective cohort study of 120 075 UK Biobank participants. *Occup Environ*
296 *Med*. 2020.
- 297 4. COVID-19 Telephone Assessment, Testing Pathway and Return to Work of
298 Symptomatic Healthcare Workers Version 9: Health Service Executive (HSE); Health
299 Protection Surveillance Centre (HPSC); 2021 [updated April 15 2021. Available from:
300 [https://www.hse.ie/eng/staff/workplace-health-and-wellbeing-unit/covid-19-](https://www.hse.ie/eng/staff/workplace-health-and-wellbeing-unit/covid-19-guidance/assessment-testing-and-return-to-work-of-symptomatic-healthcare-worker1.pdf)
301 [guidance/assessment-testing-and-return-to-work-of-symptomatic-healthcare-worker1.pdf](https://www.hse.ie/eng/staff/workplace-health-and-wellbeing-unit/covid-19-guidance/assessment-testing-and-return-to-work-of-symptomatic-healthcare-worker1.pdf).
- 302 5. Prevention, identification and management of health worker infection in the context of
303 COVID-19: World Health Organization (WHO); 2020 [updated October 30 2020. Available
304 from: <https://www.who.int/publications/i/item/10665-336265>.
- 305 6. Return to Work Criteria for Healthcare Personnel with SARS-CoV-2 Infection
306 (Interim Guidance): Centers for Disease Control and Prevention (CDC); National Center for
307 Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases; 2021 [updated
308 June 2 2021. Available from: [https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-](https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-work.html)
309 [work.html](https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-work.html).
- 310 7. Richtlijn Corona: GGZ Standaarden; 2021 [updated June 24 2021. Available from:
311 [https://www.ggzstandaarden.nl/richtlijnen/ggz-en-corona-richtlijn/richtlijn/verantwoord-](https://www.ggzstandaarden.nl/richtlijnen/ggz-en-corona-richtlijn/richtlijn/verantwoord-inzetten-van-zorg-professionals)
312 [inzetten-van-zorg-professionals](https://www.ggzstandaarden.nl/richtlijnen/ggz-en-corona-richtlijn/richtlijn/verantwoord-inzetten-van-zorg-professionals).
- 313 8. Infection prevention and control and preparedness for COVID-19 in healthcare
314 settings - sixth update: European Centre for Disease Prevention and Control (ECDC); 2021
315 [updated February 9 2021. Available from: [https://www.ecdc.europa.eu/en/publications-](https://www.ecdc.europa.eu/en/publications-data/infection-prevention-and-control-and-preparedness-covid-19-healthcare-settings)
316 [data/infection-prevention-and-control-and-preparedness-covid-19-healthcare-settings](https://www.ecdc.europa.eu/en/publications-data/infection-prevention-and-control-and-preparedness-covid-19-healthcare-settings).
- 317 9. van Kampen JJA, van de Vijver D, Fraaij PLA, et al. Duration and key determinants
318 of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-
319 19). *Nat Commun*. 2021;**12**(1):267.
- 320 10. Bullard J, Dust K, Funk D, et al. Predicting Infectious Severe Acute Respiratory
321 Syndrome Coronavirus 2 From Diagnostic Samples. *Clin Infect Dis*. 2020;**71**(10):2663-6.
- 322 11. Walsh KA, Jordan K, Clyne B, et al. SARS-CoV-2 detection, viral load and infectivity
323 over the course of an infection. *J Infect*. 2020;**81**(3):357-71.
- 324 12. Sohn Y, Jeong SJ, Chung WS, et al. Assessing Viral Shedding and Infectivity of
325 Asymptomatic or Mildly Symptomatic Patients with COVID-19 in a Later Phase. *J Clin Med*.
326 2020;**9**(9).
- 327 13. Laferl H, Kelani H, Seitz T, et al. An approach to lifting self-isolation for health care
328 workers with prolonged shedding of SARS-CoV-2 RNA. *Infection*. 2021;**49**(1):95-101.
- 329 14. Glans H, Gredmark-Russ S, Olausson M, et al. Shedding of infectious SARS-CoV-2
330 by hospitalized COVID-19 patients in relation to serum antibody responses. *BMC Infect Dis*.
331 2021;**21**(1):494.
- 332 15. Kim YI, Kim SM, Park SJ, et al. Critical role of neutralizing antibody for SARS-CoV-
333 2 reinfection and transmission. *Emerg Microbes Infect*. 2021;**10**(1):152-60.
- 334 16. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-
335 nCoV) by real-time RT-PCR. *Euro Surveill*. 2020;**25**(3).

17. Hadfield J, Megill C, Bell SM, et al. Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics*. 2018;**34**(23):4121-3.
18. Yu G, Smith DK, Huachen Zhu YG, Lam TT-Y. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution*. January 2017;**8**(1):28-36.
19. Igloi Z, Velzing J, Huisman R, et al. Clinical evaluation of the SD Biosensor SARS-CoV-2 saliva antigen rapid test with symptomatic and asymptomatic, non-hospitalized patients. *PLoS One*. 2021; **16**(12): e0260894.
20. Gniazdowski V, Morris CP, Wohl S, et al. Repeated Coronavirus Disease 2019 Molecular Testing: Correlation of Severe Acute Respiratory Syndrome Coronavirus 2 Culture With Molecular Assays and Cycle Thresholds. *Clin Infect Dis*. 2021; **73**(4): e860–e869.
21. Zhou B, Thao TTN, Hoffmann D, et al. SARS-CoV-2 spike D614G change enhances replication and transmission. *Nature*. 2021;**592**(7852):122-7.
22. Sabino EC, Buss LF, Carvalho MPS, et al. Resurgence of COVID-19 in Manaus, Brazil, despite high seroprevalence. *Lancet*. 2021;**397**(10273):452-5.
23. Sikkens JJ, Buis DTP, Peters EJG, et al. Serologic Surveillance and Phylogenetic Analysis of SARS-CoV-2 Infection Among Hospital Health Care Workers. *JAMA Netw Open*. 2021;**4**(7):e2118554.
24. Schneider S, Piening B, Nouri-Pasovsky PA, Kruger AC, Gastmeier P, Aghdassi SJS. SARS-Coronavirus-2 cases in healthcare workers may not regularly originate from patient care: lessons from a university hospital on the underestimated risk of healthcare worker to healthcare worker transmission. *Antimicrob Resist Infect Control*. 2020;**9**(1):192.
25. Koopsen J, Dekker M, Thung P, et al. Rapid reinfection with SARS-CoV-2 variant-of-concern Alpha detected in a nurse during an outbreak at a non-covid inpatient ward: lessons learned. *Antimicrob Resist Infect Control*. 2021;**10**(1):137.
26. Fafi-Kremer S, Bruel T, Madec Y, et al. Serologic responses to SARS-CoV-2 infection among hospital staff with mild disease in eastern France. *EBioMedicine*. 2020;**59**:102915.
27. Brandstetter S, Roth S, Harner S, et al. Symptoms and immunoglobulin development in hospital staff exposed to a SARS-CoV-2 outbreak. *Pediatr Allergy Immunol*. 2020;**31**(7):841-7.
28. Fill Malfertheiner S, Brandstetter S, Roth S, et al. Immune response to SARS-CoV-2 in health care workers following a COVID-19 outbreak: A prospective longitudinal study. *J Clin Virol*. 2020;**130**:104575.

TABLES

Table I. Descriptive statistics of the study cohort

Characteristic	Value
Age, median (IQR)	33 (19.0)
Female, No. (%)	54 (75.0)
Body mass index, median (IQR)	23 (6.3)
Profession, No. (%)	
Direct patient contact	44 (61.1)
Physician	10 (15.3)
Nurse	20 (27.8)
Medical intern	8 (11.1)
Clinical assistant	4 (5.6)
Other	2 (2.8)
No direct patient contact	28 (38.9)
Researcher	10 (13.9)
Pharmacy staff/assistant	5 (6.9)
Laboratory technician	2 (2.8)
Other	11 (15.3)
Comorbidities, No. (%)	
High blood pressure	3 (4.2)
Diabetes	1 (1.4)
Cardiovascular disease	1 (1.4)
Asthma	4 (5.6)
Other	4 (5.6)
Continued to work while having symptoms, No. (%)	
Yes ^a	15 (20.8)
No knowledge of regulations	0 (0.0)
Mild symptoms	12 (80.0)
Devoted symptoms to another cause	7 (40.0)
Work pressure/sense of responsibility	3 (20.0)
No	48 (66.7)
Don't Know	3 (4.2)
Unknown	6 (8.3)

^a Multiple answers were possible

Table II. Detailed symptomatology in HCWs with RT-PCR confirmed COVID-19

Symptom	Time of interview				
	Disease onset (n=72)	Day 2 (n=72)	Day 7 (n=71)	Day 14 (n=71)	Day 21 (n=71)
Respiratory symptoms					
Coughing	22 (30.6)	39 (54.9)	27 (38.0)	12 (16.9)	9 (12.7)
Pharyngitis	21 (29.2)	19 (26.8)	7 (9.9)	6 (8.5)	3 (4.2)
Dyspnoea	7 (9.7)	11 (15.5)	11 (15.5)	5 (7.0)	9 (12.7)
Rhinitis	30 (41.7)	48 (67.6)	29 (40.8)	11 (15.5)	8 (11.3)
Gastro intestinal symptoms					
Abdominal pain	4 (5.6)	7 (9.9)	3 (4.2)	2 (2.8)	0 (0.0)
Diarrhoea	7 (9.7)	8 (11.1)	2 (2.8)	1 (1.4)	1 (1.4)
Nausea	3 (4.2)	7 (9.9)	2 (2.8)	2 (2.8)	3 (4.2)
Vomiting	1 (1.4)	3 (4.2)	0 (0.0)	0 (0.0)	0 (0.0)
Anorexia	12 (16.7)	26 (36.6)	20 (28.2)	5 (7.0)	4 (5.6)
Other symptoms					
Fever	13 (18.1)	18 (25.4)	4 (5.6)	1 (1.4)	0 (0.0)
Myalgia	19 (26.4)	23 (32.4)	9 (12.7)	3 (4.2)	3 (4.2)
Headache	37 (51.4)	39 (54.9)	16 (22.5)	12 (16.9)	9 (12.7)
Fatigue	32 (44.4)	49 (69.0)	35 (49.3)	22 (31.0)	18 (25.4)
Anosmia or dysgeusia	13 (18.9)	25 (35.2)	36 (50.7)	22 (31.0)	17 (23.9)

No symptoms experienced	0 (0.0)	0 (0.0)	15 (21.1)	36 (50.7)	40 (56.3)
<i>HCWs = healthcare workers; COVID-19 = coronavirus disease 2019</i>					

FIGURE LEGENDS

Figure 1. Maximum likelihood phylogeny of SARS-CoV-2 sequences with identified potential transmission clusters.

A condensed maximum-likelihood phylogeny of SARS-CoV-2 sequences that were collected (marked with tip shapes) and a random sample of contemporaneous reference sequences (no tips) circulating within the Netherlands. Tip shapes are coloured according to the wards the HCWs (circle and square tips) and their within-department-contacts (diamond tips) were working on. The Figure zooms in on two potential transmission clusters that were found.

Table A.I. Contemporaneous SARS-CoV-2 viral genomes from the Netherlands, derived from the GISAID database

Journal Pre-proof

